

PATIENT INFORMATION
JAYPIRCA® (JAY-PIHR-KAA)
pirtobrutinib
tablets

What is JAYPIRCA?

JAYPIRCA is a prescription medicine used to treat adults with:

- mantle cell lymphoma (MCL) that has come back or did not respond to previous treatment and who have already received at least 2 treatments for their cancer, including a Bruton tyrosine kinase (BTK) inhibitor medicine.
- chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) who have already received at least 2 treatments for their cancer, including a BTK inhibitor medicine and a B-cell lymphoma 2 (BCL-2) inhibitor medicine.

It is not known if JAYPIRCA is safe and effective in children.

Before taking JAYPIRCA, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection or have been advised that you are at increased risk of infection
- have had recent surgery or plan to have surgery. Your healthcare provider may stop JAYPIRCA for any planned medical, surgical, or dental procedure.
- have bleeding problems or are taking a blood thinner medicine
- have or had heart rhythm problems
- have high blood pressure
- have a history of other cancers including skin cancer
- have kidney problems
- are pregnant or plan to become pregnant. JAYPIRCA can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before starting treatment with JAYPIRCA.
- You should use effective birth control (contraception) during treatment with JAYPIRCA and for 1 week after your last dose of JAYPIRCA.
- Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with JAYPIRCA.
- are breastfeeding or plan to breastfeed. It is not known if JAYPIRCA passes into your breast milk. Do not breastfeed during treatment with JAYPIRCA and for 1 week after your last dose of JAYPIRCA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking JAYPIRCA with certain other medicines may affect how JAYPIRCA or the other medicines work and can cause side effects.

How should I take JAYPIRCA?

- Take JAYPIRCA exactly as your healthcare provider tells you.
- Do not change your dose or stop taking JAYPIRCA unless your healthcare provider tells you to do so.
- Take JAYPIRCA tablets 1 time each day at about the same time each day.
- Take JAYPIRCA with or without food.
- Swallow JAYPIRCA tablets whole with water. Do not cut, crush, or chew the tablets.
- If you miss a dose of JAYPIRCA, take it as soon as you remember on the same day. If it has been more than 12 hours from the time you usually take JAYPIRCA, skip the missed dose, and take your next dose on the next day at your usual time.

What are the possible side effects of JAYPIRCA?

JAYPIRCA can cause serious side effects, including:

- **Infections** can happen during treatment with JAYPIRCA. These infections can be serious and may lead to death. Your healthcare provider may prescribe vaccines and certain medicines if you have an increased risk of getting infections. Tell your healthcare provider right away if you develop fever, chills, weakness, flu-like symptoms, or any other signs of infection during treatment with JAYPIRCA.
- **Bleeding problems (hemorrhage)** can happen during treatment with JAYPIRCA and can be serious and may lead to death. Your risk of severe bleeding may increase if you are also taking a blood thinner medicine. Tell your healthcare provider if you develop any signs or symptoms of bleeding, including:
 - blood in your stools or black stools (looks like tar)
 - pink or brown urine
 - unexpected bleeding, or bleeding that is severe or you cannot control
 - vomit blood or vomit blood that looks like coffee grinds
 - cough up blood or blood clots
 - increased bruising
 - dizziness
 - weakness
 - confusion
 - changes in your speech
 - headache that lasts a long time

- **Decrease in blood cell counts.** Decrease in blood cell counts (white blood cells, platelets, and red blood cells) are common with JAYPIRCA, but can also be severe. This may increase your risk of infection, bleeding, and anemia. Your healthcare provider should do blood tests to check your blood counts regularly during treatment with JAYPIRCA.
- **Heart rhythm problems.** Heart rhythm problems including atrial fibrillation and atrial flutter have happened in people treated with JAYPIRCA. Your risk for heart rhythm problems may be increased if you have high blood pressure or have had heart rhythm problems in the past. Tell your healthcare provider if you develop any of the following signs or symptoms:
 - fast or irregular heartbeat (palpitations)
 - chest discomfort
 - dizziness
 - shortness of breath
 - fainting
- **Second primary cancers.** New cancers have happened in people during treatment with JAYPIRCA, including cancers of the skin or other organs. Your healthcare provider will check you for other cancers during treatment with JAYPIRCA. Use sun protection when you are outside in sunlight.

Your healthcare provider may decrease your dose, temporarily stop, or permanently stop treatment with JAYPIRCA if you develop severe side effects.

The most common side effects of JAYPIRCA include:

- tiredness
- muscle, joint, and bone pain
- diarrhea
- COVID-19
- bruising
- cough

These are not all the possible side effects of JAYPIRCA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store JAYPIRCA?

- Store JAYPIRCA at room temperature between 68°F to 77°F (20°C to 25°C).
- JAYPIRCA comes in a bottle with a child-resistant cap.

Keep JAYPIRCA and all medicines out of the reach of children.

General information about the safe and effective use of JAYPIRCA.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use JAYPIRCA for a condition for which it was not prescribed. Do not give JAYPIRCA to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about JAYPIRCA that is written for health professionals.

What are the ingredients in JAYPIRCA?

Active ingredient: pirtobrutinib

Inactive ingredients: croscarmellose sodium, hypromellose acetate succinate, lactose monohydrate, magnesium stearate, microcrystalline cellulose, and silicon dioxide. The tablet film coating contains FD&C Blue #2, hypromellose, titanium dioxide, and triacetin.

Marketed by Lilly USA, LLC, Indianapolis, IN 46285, USA

JAYPIRCA is a trademark of Eli Lilly and Company.
Copyright © 2023, Eli Lilly and Company. All rights reserved.
JAY-0003-PPI-20231215

For more information, go to www.jaypirca.com or call 1-800-LillyRx (1-800-545-5979).

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 12/2023