

## INSTRUCTIONS FOR USE

**PrOMVOH™ (ahm-VOH)  
mirikizumab injection  
for subcutaneous use  
Prefilled Syringe**



**This Instructions for Use contains information on how to inject OMVOH.**

**For subcutaneous injection only.**

**Before you use OMVOH prefilled syringes, read and carefully follow all the step-by-step instructions. Two injections are required for a full dose.**

### **Important information you need to know before injecting OMVOH**


- Your healthcare provider should show you how to prepare and inject OMVOH using the prefilled syringe. **Do not** inject yourself or someone else until you have been shown how to inject OMVOH.
- Keep this Instructions for Use and read it as needed.
- 2 OMVOH injections are required for a full dose.
- Inject 1 OMVOH prefilled syringe followed right away by the other OMVOH prefilled syringe.
- Each OMVOH prefilled syringe is for **one-time use only**. **Do not** share or reuse your OMVOH prefilled syringe. You may give or get an infection.
- Your healthcare provider may help you decide where on your body to inject your dose. You can also read the **Choose your injection site** section of these instructions to help you choose which area can work best for you.
- If you have vision problems, **do not** use OMVOH prefilled syringe without help from a caregiver.
- See **Storing OMVOH Prefilled Syringes** for important storage information.

## INSTRUCTIONS FOR USE

Before you use OMVOH prefilled syringes, read and carefully follow all the step-by-step instructions.

### Parts of the OMVOH Prefilled Syringe

Top



Bottom

100 mg/mL + 100 mg/mL = 1 Full Dose

#### **IMPORTANT:**

- 2 OMVOH injections are required for a full dose.
- Inject 1 OMVOH prefilled syringe followed right away by the other OMVOH prefilled syringe.

## Preparing to inject OMVOH

### Take the Prefilled Syringes from the refrigerator

Take 2 OMVOH prefilled syringes from the refrigerator.

**Leave the needle caps on until you are ready to inject.**

Leave the prefilled syringes at room temperature for 30 minutes before injecting.

**Do not** microwave the prefilled syringes, or run hot water over them, or leave them in direct sunlight.

**Do not** use the prefilled syringes if the medicine is frozen.


**Do not** shake the prefilled syringes.

### Gather supplies

Supplies:

- 2 alcohol wipes
- 2 cotton balls or pieces of gauze
- 1 sharps container (see **Disposing of OMVOH Prefilled Syringes**)

### Inspect the Prefilled Syringes and the medicine



Make sure you have the right medicine. The medicine inside should be clear. It may be colourless to slightly yellow.

**Do not** use the prefilled syringes and throw away (dispose of) as directed by your healthcare provider or pharmacist if:


- one or both prefilled syringes look damaged
- the medicine is cloudy, is discoloured, or has particles
- the expiration date printed on the label has passed
- the medicine is frozen

### Prepare for injection

Wash your hands with soap and water before you inject OMVOH.

### Choose your injection site

Your healthcare provider can help you choose the injection site that is best for you.




- **You or another person** may inject the medicine into your stomach area (abdomen). **Do not** inject within 2 inches of the belly button (navel).
- **You or another person** may inject the medicine in the front of your thighs. This area should be at least 2 inches above the knee and 2 inches below the groin.
- **Another person** may give you the injection in the back of your upper arm.
- **Do not** inject in the exact same spot every time. For example, if your first injection was in your abdomen, your second injection — to complete a full dose — could be in another spot in your abdomen.
- **Do not** inject into areas where the skin is tender, bruised, red, or hard.

**Clean the injection sites with an alcohol wipe. Let the injection sites dry before you inject the medicine.**

## Injecting OMVOH


### 1 Uncap

- **Leave the needle cap on until you are ready to inject.**
- Pull the needle cap off and throw it away in your household trash.
- **Do not** put the needle cap back on – you could damage the needle or stick yourself by accident.
- **Do not** touch the needle.



### 2 Insert


- Gently pinch and hold a fold of skin where you will inject.



- Insert the needle at a 45-degree angle.


### 3 Inject

- Slowly push on the thumb pad to push the plunger all the way in until all the medicine is injected.



- The gray syringe plunger should be pushed all the way to the needle end of the syringe.

- You should see the blue plunger rod show through the syringe body when the injection is complete as shown.
- Remove the needle from your skin and gently let go of your skin.
- If you have bleeding at the injection site, press a cotton ball or gauze over the injection site.
- **Do not** rub the injection site, as this may cause bruising.
- **Do not** put the needle cap back on the prefilled syringe.



**Blue Plunger Rod**

**Gray Syringe Plunger**

**2 injections are required for a full dose. Inject one prefilled syringe immediately followed by the other prefilled syringe.**

### Disposing of OMVOH Prefilled Syringes

#### Throw away the used prefilled syringes

- Put the used OMVOH prefilled syringes in a sharps disposal container right away after use. **Do not** throw away (dispose of) the OMVOH prefilled syringes in your household trash.



- If you do not have a sharps disposal container, you may use a household container that is:
  - made of a heavy-duty plastic,
  - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
  - upright and stable during use,
  - leak-resistant, and
  - properly labeled to warn of hazardous waste inside the container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be local laws about how you should throw away needles and syringes.
- **Do not** recycle your used sharps disposal container.

## Commonly Asked Questions

**Q. What if I let my OMVOH prefilled syringe warm up for longer than 30 minutes before injecting?**

**A.** Your prefilled syringe can stay at room temperature up to 30°C for up to 2 weeks.

**Q. What if I see air bubbles in the OMVOH prefilled syringe?**

**A.** It is normal to have air bubbles in the prefilled syringe. They will not harm you or affect your dose.

**Q. What if there is a drop of liquid on the tip of the needle when I remove the needle cap?**

**A.** It is okay to see a drop of liquid on the tip of the needle. This will not harm you or affect your dose.

**Q. What if I cannot push in the plunger?**

**A.** If the plunger is stuck or damaged:

- **Do not** continue to use the syringe
- Remove the needle from your skin
- Dispose of the syringe and get a new one

**Q. What if there is a drop of liquid or blood on my skin after my injection?**

**A.** This is normal. Press a cotton ball or gauze over the injection site. **Do not** rub the injection site.

**Q. How can I tell if my injection is complete?**

**A.** When your injection is complete:

- The blue plunger rod should show through the body of the syringe.
- The gray syringe plunger should be pushed all the way to the needle end of the syringe.

**If you have more questions about how to use the OMVOH prefilled syringe:**

- Call your healthcare provider or
- Call Lilly at 1-888-545-5972
- Visit [www.lilly.ca](http://www.lilly.ca)

## Storing OMVOH Prefilled Syringes

- Store your prefilled syringes in the refrigerator between 2°C to 8°C.
- Your prefilled syringes may be stored at room temperature for up to 2 weeks. **Do not** store above 30°C. Throw away (dispose of) OMVOH if not used within 2 weeks at room temperature.
- **Do not** freeze your prefilled syringes.
- Store your prefilled syringes in the original to protect from light until use.
- **Do not** microwave your prefilled syringes, or run hot water over them, or leave them in direct sunlight.

- **Do not** shake your prefilled syringes.
- Throw away (dispose of) your prefilled syringes if any of the above conditions are not followed.
- **Keep your prefilled syringes and all medicines out of the sight and reach of children.**

**Read the Patient Medication Information for OMVOH inside this box to learn about your medicine.**

Eli Lilly Canada Inc., P.O. Box 73, Toronto, Ontario, M5X 1B1

OMVOH™ is a trademark of Eli Lilly and Company.

Copyright © 2023, Eli Lilly and Company. All rights reserved.

Document Issued: July 20, 2023

OMV-0001-CA-PFS-IFU-20230720